

Back to bright

Opalescence™ Endo Whitening

- 35% hydrogen peroxide
- Specifically formulated to whiten endodontically treated, discolored, non-vital teeth
- Designed for the “walking bleach” technique
- Convenient, premixed syringe delivery



Non-Vital “Walking Bleach”



Before



Before



After



After

Opalescence™ Endo Whitening

MUST BE REFRIGERATED

1270 - Opalescence Endo Syringe Kit

2 x 1.2 ml Opalescence Endo syringes
20 x Black Mini tips



MUST BE REFRIGERATED

1323 - Opalescence Endo Syringe 2pk

1.2 ml syringes



Procedure

Courtesy of Dr. Carlos Ramos



1. Completely remove all the restorative and root sealing material from the coronal pulp chamber and 2-3 mm below healthy gingiva. Place a 2 mm thick conventional glass ionomer or a resin modified glass ionomer to seal the endodontically treated canal. Verify set of material before proceeding.



2. Apply a layer of Opalescence Endo gel to the chamber.



3. Insert a cotton pellet into the chamber.



4. Place mixed UltraTemp™ Regular filling material directly to site.



5. Wipe away excess with a wet cotton ball or gauze before it sets.



Finished. Repeat every 1-5 days until desired results are achieved.

