

Opalescence[™]
tooth whitening systems



Every Smile Brightened!




Opalescence™ Tooth Whitening Reference Guide




Opalescence™ PF

 Custom Whitening Tray
(Requires Impression)

Opalescence™ PF 10%

 **Wear Time:**
8–10 hours per day


 **Flavors:**
Mint, Melon, and Regular


Active Ingredient:
10% carbamide peroxide

Contains:
Potassium nitrate and fluoride

Indications:
For patients with existing sensitivity; can be worn day or night

Opalescence™ PF 15% and 20%

 **Wear Time:**
15%: 4–6; 20%: 2–4 hours per day


 **Flavors:**
Mint, Melon, and Regular


Active Ingredient:
15% and 20% carbamide peroxide

Contains:
Potassium nitrate and fluoride

Indications:
For faster whitening; can be worn during the day

Opalescence™ PF 35%

 **Wear Time:**
30–60 minutes a day


 **Flavors:**
Mint, Melon, and Regular

Active Ingredient:
35% carbamide peroxide

Contains:
Potassium nitrate and fluoride

Indications:
For shorter wear time and touch-ups

Opalescence™ PF 45%

 **Wear Time:**
15–30 minutes

 **Flavors:**
Mint

Active Ingredient:
45% carbamide peroxide

Contains:
Potassium nitrate and fluoride

Indications:
For shorter wear times




Opalescence Go™

 UltraFit™ Ready-To-Go Tray

Opalescence Go™ 10%

 **Wear Time:**
30–60 minutes per day

 **Flavors:**
Mint and Melon

Active Ingredient:
10% hydrogen peroxide

Contains:
Potassium nitrate and fluoride

Indications:
For fast whitening ready-to-go; an alternative to OTC products

Opalescence Go™ 15%

 **Wear Time:**
15–20 minutes per day

 **Flavors:**
Mint

Active Ingredient:
15% hydrogen peroxide

Contains:
Potassium nitrate and fluoride


Indications:
For fast whitening ready-to-go; an alternative to OTC products



Opalescence™ Endo

 Applied In-Office

Opalescence™ Endo

 **Wear Time:**
1–5 days per application

Active Ingredient:
35% hydrogen peroxide


Indications:
Dentist administered for internal whitening of non-vital teeth



Opalustre™

 Applied In-Office

Opalustre™ Microabrasion Slurry

 **Wear Time:**
1 or more doctor-administered applications as needed

Active Ingredient:
6.6% hydrochloric acid


Indications:
Dentist-administered chairside treatment; removes superficial white and brown stains



Opalescence™ Boost™

 Applied In-Office

Opalescence™ Boost™

 **Wear Time:**
Two to three 20-minute professionally administered treatments per visit (do not exceed three per visit)

Active Ingredient:
40% hydrogen peroxide

Contains:
Potassium nitrate and fluoride

Indications:
Dentist-administered, fast chairside treatment



UltraEZ™

 UltraFit™ Ready-To-Go Tray

 Custom Whitening Tray
(Requires Impression)

UltraEZ™

 **Wear Time:**
15–60 minutes per day

 **Flavors:**
Regular

Active Ingredient:
3% potassium nitrate, .25% neutral NaF

Indications:
Treats sensitivity