

# **Break the Tension**

ChlorCid 3% sodium hypochlorite solution with a deeper cleaning power





## ChlorCid<sup>™</sup> Surf

**ChlorCid™Surf** 3% sodium hypochlorite solution includes surfactant to modify surface tension. This allows irrigant to flow into accessory canals of the root canal system to give a deeper clean.

### ChlorCid<sup>™</sup>Surf

Sodium hypochlorite (NaOCL) solutions are the most common endodontic irrigant due to their cleaning and tissue dissolution properties. The addition of surfactant increases the ability of the NaOCL to penetrate the root canal isthmuses, allowing for greater cleaning action on dentinal walls.

> 4613 - ChlorCid Surf Bottle 1pk 480 ml bottle

97 - ChlorCid Surf IndiSpense Syringe 1pk 30 ml syringe



Sodium hypochlorite with surfactant used with ultrasonic movement has been shown to be the most effective method for soft tissue dissolution.1



### Other irrigants available from Ultradent

3% Sodium Hypochlorite

4612 - ChlorCid Bottle 1pk 480 ml bottle



1467 - ChlorCid V IndiSpense Syringe 1pk 30 ml syringe



**EDTA 18% for smear layer removal** 

4616 - EDTA Bottle 1pk 480 ml bottle



162 - EDTA IndiSpense Syringe 1pk 30 ml syringe



#### 2% Chlorhexidine Antibacterial Solution

4614 - Consepsis Bottle 1pk 480 ml bottle



687 - Consepsis IndiSpense Syringe 1pk 30 ml syringe



2210 - Consepsis V IndiSpense Syringe 1pk 30 ml syringe



1. Estevez R, Conde AJ, de Pablo OV, de la Torre F, Rossi-Fedele G, MclinDent, Cisneros R. Effect of passive ultrasonic activation on organic tissue dissolution from simulated grooves in root canals using sodium hypochlorite with or without surfactants and EDTA. JOE. 2017;43(7):1161-1165.





