

Opalescence™
tooth whitening systems



Every Smile Brightened!




Opalescence™ Tooth Whitening Reference Guide




Opalescence™ PF whitening

 Custom Whitening Tray
(Requires Impression)

Opalescence™ PF 10%

 **Wear Time:**
8–10 hours per day


 **Flavors:**
Mint, Melon, and Regular


Whitening Agent:
10% carbamide peroxide

Contains:
Potassium nitrate and fluoride

Use:
For patients with sensitivity concerns; can be worn day or night

Opalescence™ PF 15% and 20%

 **Wear Time:**
15%: 4–6; 20%: 2–4 hours per day

 **Flavors:**
Mint, Melon, and Regular


Whitening Agent:
15% and 20% carbamide peroxide

Contains:
Potassium nitrate and fluoride

Use:
For faster whitening; daytime wear recommended

Opalescence™ PF 35%

 **Wear Time:**
30–60 minutes a day


 **Flavors:**
Mint, Melon, and Regular


Whitening Agent:
35% carbamide peroxide

Contains:
Potassium nitrate and fluoride

Use:
For shorter wear time and touch-ups

Opalescence™ Quick PF 45%

 **Wear Time:**
15–30 minutes

 **Flavors:**
Mint

Whitening Agent:
45% carbamide peroxide

Contains:
Potassium nitrate and fluoride

Use:
For shorter wear times




Opalescence Go™ whitening

 UltraFit™ Ready-To-Go Tray

Opalescence Go™ 10%

 **Wear Time:**
30–60 minutes per day


 **Flavors:**
Mint and Melon

Whitening Agent:
10% hydrogen peroxide

Contains:
Potassium nitrate and fluoride

Use:
For fast whitening ready-to-go; an alternative to OTC products

Opalescence Go™ 15%

 **Wear Time:**
15–20 minutes per day

 **Flavors:**
Mint

Whitening Agent:
15% hydrogen peroxide

Contains:
Potassium nitrate and fluoride


Use:
For fast whitening ready-to-go; an alternative to OTC products



Opalescence™ Endo whitening

 Applied In-Office

Opalescence™ Endo

 **Wear Time:**
1–5 days per application

Whitening Agent:
35% hydrogen peroxide


Use:
Dentist-administered for internal whitening of endodontically treated teeth



Opalustre™ slurry

 Applied In-Office

Opalustre™ Microabrasion Slurry

 **Application:**
1 or more doctor-administered applications as needed

Removal Agent:
6.6% hydrochloric acid silicone carbide


Use:
Dentist-administered chairside treatment; permanently removes superficial enamel imperfections



Opalescence™ Boost™ whitening

 Applied In-Office

Opalescence™ Boost™

 **Application:**
Two to three 20-minute professionally administered applications per visit (do not exceed three per visit)

Whitening Agent:
40% hydrogen peroxide

Contains:
Potassium nitrate and fluoride

Use:
Dentist-administered, fast chairside treatment




UltraEZ™ desensitizing gel

 UltraFit™ Ready-To-Go Tray

 Custom Whitening Tray
(Requires Impression)

UltraEZ™

 **Wear Time:**
15–60 minutes per day

 **Flavors:**
Regular

Desensitizing Agent:
3% potassium nitrate, .25% neutral NaF

Use:
Treats sensitivity