

Opalescence™

tooth whitening systems



Every Smile Brightened!



Opalescence™ Tooth Whitening Reference Guide



Opalescence™ PF whitening

 Custom Whitening Tray
(Requires Impression)

Opalescence™ 10%



Flavors:
Mint and
Regular



Wear Time:
8-10 hours
per day

Whitening Agent:

10% carbamide peroxide

Use:

For patients with sensitivity concerns; can be worn day or night

Opalescence™ PF 10%



Flavors:
Mint, Melon,
and Regular



Wear Time:
8-10 hours
per day

Whitening Agent:

10% carbamide peroxide

Contains:

Potassium nitrate and fluoride

Use:

For patients with sensitivity concerns; can be worn day or night

Opalescence™ PF 15% and 20%



Flavors:
Mint, Melon,
and Regular



Wear Time:
15%: 4-6; 20%:
2-4 hours per day

Whitening Agent:

15% and 20% carbamide peroxide

Contains:

Potassium nitrate and fluoride

Use:

For faster whitening; daytime wear recommended

Opalescence™ PF 35%



Flavors:
Mint, Melon,
and Regular



Wear Time:
30-60 minutes
per day

Whitening Agent:

35% carbamide peroxide

Contains:

Potassium nitrate and fluoride

Use:

For shorter wear time and touch-ups

Opalescence™ Quick PF 45%



Flavors:
Mint



Wear Time:
15-30 minutes

Whitening Agent:

45% carbamide peroxide

Contains:

Potassium nitrate and fluoride

Use:

For shorter wear times



Opalescence Go™ whitening

 UltraFit™ Ready-To-Go Tray

Opalescence Go™ 10%



Wear Time:
30-60 minutes per day



Flavors:
Mint and Melon

Whitening Agent:

10% hydrogen peroxide

Contains:

Potassium nitrate and fluoride

Use:

For fast whitening ready-to-go; an alternative to OTC products

Opalescence Go™ 15%



Wear Time:
15-20 minutes per day



Flavors:
Mint

Whitening Agent:

15% hydrogen peroxide

Contains:

Potassium nitrate and fluoride

Use:

For fast whitening ready-to-go; an alternative to OTC products



Opalescence™ Endo whitening

 Applied In-Office

Opalescence™ Endo



Wear Time:
1-5 days per application

Whitening Agent:

35% hydrogen peroxide

Use:

Dentist-administered for internal whitening of endodontically treated teeth



Opalustre™ slurry

 Applied In-Office

Opalustre™ Microabrasion Slurry



Application:
1 or more doctor-administered
applications as needed

Removal Agent:

6.6% hydrochloric acid silicone carbide

Use:

Dentist-administered chairside treatment; permanently removes superficial enamel imperfections



Opalescence™ Boost™ whitening

 Applied In-Office

Opalescence™ Boost™



Application:
Two to three 20-minute professionally
administered applications per visit (do
not exceed three per visit)

Whitening Agent:

40% hydrogen peroxide

Contains:

Potassium nitrate and fluoride

Use:

Dentist-administered, fast chairside treatment



UltraEZ™ desensitizing gel

 UltraFit™ Ready-To-Go Tray

 Custom Whitening Tray
(Requires Impression)

UltraEZ™



Wear Time:
15-60 minutes per day



Flavors:
Regular

Desensitizing Agent:

3% potassium nitrate, .25% neutral NaF

Use:

Treats sensitivity