

BENYLIN WET COUGH MENTHOL

SCHEDULING STATUS:

S0

PROPRIETARY NAME (AND DOSAGE FORM):

Benylin Wet Cough Menthol (Syrup)

COMPOSITION:

Quantity per 10 ml:

Guaifenesin	200 mg
Alcohol	9,5% v/v
Sugar	2,0 g
Glucose	5,0 ml
Preservative:	
Sodium Benzoate	0,2% m/v

PHARMACOLOGICAL CLASSIFICATION:

A: 10.1 Antitussives and Expectorants.

PHARMACOLOGICAL ACTION:

Guaifenesin has expectorant properties.

INDICATIONS:

Alleviation of cough.

CONTRA-INDICATIONS:

Allergy to any of the ingredients. Pregnancy and lactation.

Guaifenesin is considered unsafe in patients with acute porphyria.

DOSAGE AND DIRECTIONS FOR USE:

Adults: 10 ml - 20 ml (two - four medicine measures) every 4 hours.

If symptoms persist a doctor should be consulted.

SIDE-EFFECTS AND SPECIAL PRECAUTIONS:

Gastro-intestinal discomfort and drowsiness may occur.

Guaifenesin may interfere with diagnostic measurements of urinary 5-hydroxy-indoleacetic acid or vanillylmandelic acid.

KNOWN SYMPTOMS OF OVERDOSAGE AND PARTICULARS OF ITS TREATMENT:

In large doses, Guaifenesin will cause drowsiness, nausea and vomiting.

Treatment is symptomatic and supportive.

IDENTIFICATION:

A reddish-brown, syrupy liquid with a strong peppermint-caramel odour and taste.

PRESENTATION:

Bottles of 50 ml and 100 ml, with a plastic measuring cup.

STORAGE INSTRUCTIONS:

Store in a cool place (at or below 25°C).

KEEP OUT OF REACH OF CHILDREN.

REGISTRATION NUMBER:

28/10.1/0374

**NAME AND BUSINESS ADDRESS OF THE HOLDER OF THE CERTIFICATE OF
REGISTRATION:**

Johnson & Johnson (Pty) Ltd.

241 Main Road

RETREAT

7945

South Africa

DATE OF PUBLICATION OF THIS PACKAGE INSERT:

30 March 1994

EXPORT REGISTRATION DETAILS

Botswana: B9326080

Ghana: FDB/SD.07-1023

Namibia: 05/10.1/0286

Zambia: 082/039