

courte. L'incidence de l'intubation difficile chez l'enfant âgé de plus d'un an est très rare et souvent associée à un syndrome polymalformatif. Dans ce cas, le recours à un vidéolaryngoscope est recommandé en première intention associé à un algorithme adapté et des aides cognitives sur la gestion de l'intubation difficile. Les indications de fibroscopie vigile et de trachéotomie sont extrêmement rares chez l'enfant.

Références

- [1] Adewale L. Anatomy and assessment of the pediatric airway. *Pediatr Anesth* 2009;19:1–8.
- [2] Wilton N, Hack H. Developmental anatomy of the airway. *Anaesth Intensive Care Med* 2019;20:29–34.
- [3] Diemunsch P, Langeron O, Richard M, Lenfant F. Prédiction et définition de la ventilation au masque difficile et de l'intubation difficile. *Ann Fr Anesth Reanim* 2008;27:3–14.
- [4] Heinrich S, Birkholz T, Ihmsen H, Irouschek A, Ackermann A, Schmidt J. Incidence and predictors of difficult laryngoscopy in 11.219 pediatric anesthesia procedures: Erlangen pediatric laryngoscopy. *Pediatr Anesth* 2012;22:729–36.
- [5] Engelhardt T, Virag K, Veyckemans F, Habre W. Airway management in paediatric anaesthesia in Europe – insights from APRICOT (Anaesthesia Practice In Children Observational Trial): A prospective multicentre observational study in 261 hospitals in Europe. *Br J Anaesth* 2018;121:66–75.
- [6] Raj D, Luginbuehl I. Managing the difficult airway in the syndromic child. *Contin Educ Anaesth Crit Care Pain* 2015;15:7–13.
- [7] Dupeyrat A, Dubreuil M, Ecoffey C. Preoxygenation in children. *Anesth Analg* 1994;79(5):1027.
- [8] Morrison J, Collier E, Friesen R, Logan L. Preoxygenation before laryngoscopy in children: How long is enough? *Pediatr Anesth* 1998;8:293–8.
- [9] Dadure C, Sabourdin N, Veyckemans F, Babre F, Bourdaud N, Dahmani S, et al. Gestion des voies aériennes de l'enfant. *Anesth Reanim* 2019;5:408–26.
- [10] Humphreys S, Lee-Archer P, Reyne G, Long D, Williams T, Schibler A. Transnasal humidified rapid-insufflation ventilatory exchange (THRIVE) in children: A randomized controlled trial. *Br J Anaesth* 2017;118:232–8.
- [11] Riva T, Pedersen TH, Seiler S, Kasper N, Theiler L, Greif R, et al. Transnasal humidified rapid insufflation ventilatory exchange for oxygenation of children during apnoea: A prospective randomised controlled trial. *Br J Anaesth* 2018;120:592–9.
- [12] Black AE, Flynn PER, Smith HL, Thomas ML, Wilkinson KA. Development of a guideline for the management of the unanticipated difficult airway in pediatric practice. *Pediatr Anesth* 2015;25:346–62.
- [13] Kemper M, Imach S, Buehler PK, Thomas J, Dave M, Weiss M. Tube tip and cuff position using different strategies for placement of currently available paediatric tracheal tubes. *Br J Anaesth* 2018;121:490–5.
- [14] Shibasaki M, Nakajima Y, Ishii S, Shimizu F, Shime N, Sessler DI. Prediction of pediatric endotracheal tube size by ultrasonography. *Anesthesiology* 2010;113:819–24.
- [15] Altun D, Orhan-Sungur M, Ali A, Özkan-Seyhan T, Sivrikoz N, Çamcı E. The role of ultrasound in appropriate endotracheal tube size selection in pediatric patients. In: Von Ungern-Sternberg B, editor. *Pediatr Anesth*; 2017. p. 1015–20. 27.
- [16] Abdelgadir IS, Phillips RS, Singh D, Moncreiff MP, Lumsden JL. Videolaryngoscopy versus direct laryngoscopy for tracheal intubation in children (excluding neonates). *Cochrane Database Syst Rev* 2017;5. CD011413.
- [17] De Carvalho CC, Regueira SLPA, Souza ABS, Medeiros LMLF, Manoel MBS, Da Silva DM, et al. Videolaryngoscopes versus direct laryngoscopes in children: Ranking systematic review with network meta-analyses of randomized clinical trials. *Pediatr Anesth* 2022;32:1000–14.